


GAZETA MATEMATICĂ

PUBLICAȚIE LUNARĂ PENTRU TINERET

7-8-9



ISSN 1584-9333

Generalizarea 1. Dacă $f, g : [0, 1] \rightarrow \mathbb{R}$ sunt funcții integrabile Riemann, atunci pentru orice sisteme de puncte intermediare ξ_1, ξ_2 asociate diviziunii $\Delta_n = \left(0 < \frac{1}{n} < \frac{2}{n} < \dots < \frac{n-1}{n} < 1\right)$, $\xi_1^k, \xi_2^k \in \left[\frac{k}{n}, \frac{k+1}{n}\right]$, $k = \overline{0, n-1}$, avem:
$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=0}^{n-1} f(\xi_1^k) g(\xi_2^k) = \int_0^1 f(x)g(x)dx.$$

Observație. În mod evident, se putea generaliza chiar pentru un șir oarecare de diviziuni cu norma tinzând la 0, dar, pentru a simplifica prezentarea, am preferat utilizarea șirului de diviziuni echidistante.

Generalizarea 2. Prin inducție, putem extinde rezultatul pentru p funcții și p sisteme de puncte intermediare, cu $p \in \mathbb{N}$, $p \geq 2$.

BIBLIOGRAFIE

- [1] Árpád Bényi, Maria Bobeș, *Asupra unei probleme de olimpiadă*, G. M.-B nr. 10/2008.
- [2] Nicu Boboc, *Analiză Matematică*, Editura Universității București, 1999.

Nota redacției. Autorii notei de față au soluționat, independent unul de celălalt, problema în cauză. Soluțiile fiind practic identice, redacția a optat pentru varianta celui de-al doilea autor, fiind mai adecvată manualelor românești în vigoare, deși, primul material primit la redacție a fost cel al primului autor.

ON A HOMOGENEOUS INEQUALITY GIVEN AT THE 2004 J.B.M.O.

VASILE BERINDE¹⁾

Abstract. Starting from a homogeneous JBMO inequality, other interesting related results are obtained.

Keywords: elementary inequality

MSC : 26D05

1. Introduction

The first problem given to the Junior Balkan Mathematical Olympiad in 2004, proposed by Albania, see [2], [3], was the following:

Problem 1. *Prove that the inequality:*

$$\frac{x+y}{x^2-xy+y^2} \leq \frac{2\sqrt{2}}{\sqrt{x^2+y^2}}, \quad (1)$$

holds for all real numbers x and y , not both equal to 0.

In the very recent book of S. Bilchev [2], two solutions of this problem are given. We present here both of them, in view of some comments and

¹⁾ Department of Mathematics and Computer Science North University of Baia Mare,
E-mail: vberinde@ubm.ro

developments¹⁾.

First Solution. If $x + y \leq 0$, the inequality obviously holds (the left hand side is negative or zero, while the right hand side is positive).

It is also easy to check that for $x = 0$ or $y = 0$, the inequality in (1) is strict.

Consider, therefore, only the case $x + y > 0$ and $x \neq 0, y \neq 0$. Then (1) can be equivalently written under the form:

$$(x + y) \cdot \frac{x^2 + y^2}{2} \leq \sqrt{2(x^2 + y^2)} \cdot (x^2 - xy + y^2). \quad (2)$$

But

$$x + y \leq \sqrt{2(x^2 + y^2)} \quad (\Leftrightarrow (x - y)^2 \geq 0) \quad (3)$$

and

$$\frac{x^2 + y^2}{2} \leq x^2 - xy + y^2 \quad (\Leftrightarrow (x - y)^2 \geq 0). \quad (4)$$

Now, in view of the fact that the numbers in both sides of (3) and (4) are positive, by multiplying (3) and (4) side by side we get exactly (2).

Equality in (1) holds if and only if equality holds in (3) and (4), that is, if $x = y$.

Second Solution. Similarly to the first solution, consider only the case $x + y > 0$ and $x \neq 0, y \neq 0$. By denoting $S = x^2 + y^2$ and $P = xy$, the inequality (1) can be equivalently written, after squaring both sides, as:

$$(S + 2P) \cdot S \leq 8(S - P)^2$$

which reduces to $7S^2 - 18SP + 8P^2 \geq 0 \Leftrightarrow (S - 2P)(7S - 4P) \geq 0$. But $S - 2P = (x - y)^2 \geq 0$, with equality if and only if $x = y$, and

$$7S - 4P = 7x^2 - 4xy + 7y^2 = y^2 \cdot \left[7\left(\frac{x}{y}\right)^2 - 4\frac{x}{y} + 7 \right] > 0,$$

for all $x, y \in \mathbb{R}, x \neq 0, y \neq 0$.

2. Other similar inequalities

Now let us have a close look on the main argument in the second solution presented above.

Basically, the key tool in proving inequality (1) was to reduce it to an inequality of the form:

$$(S - 2P)(aS + bP) \geq 0, \quad (5)$$

where a and b were some constants for which $aS + bP > 0$, for all $x, y \in \mathbb{R}, x \neq 0, y \neq 0$. As $\operatorname{sgn}(aS + bP) = \operatorname{sgn}(at^2 + bt + a)$, $t \in \mathbb{R}$, all that is needed in order to have (5) satisfied is to have: $at^2 + bt + a > 0, \forall t \in \mathbb{R}$. But, in view of the properties of the quadratic function, this happens if:

¹⁾Dedicated to the memory of my friend Professor *Svetoslav Jordanov Bilchev* (1946-2010), former Dean of Faculty of Education and Head of Department of Algebra and Geometry, University of Rousse, Bulgaria

1) $a > 0$ and 2) $\Delta = b^2 - 4a^2 < 0$,
that is, if $a > 0$ and $b \in (-2a, 2a)$. In that case (5) can be equivalently written as:

$$(a+1)(S-P)^2 \geq (a+2b+1)P^2 + S^2 - (b+2)SP. \quad (6)$$

Now, in order to get the factor S in (6), we must have:

$$a+2b+1=0, \quad (7)$$

which was satisfied in the original case of Problem 1, when we have had $a=7$ and $b=-4$.

It is easy to see that if we would like to get:

$$S^2 - (b+2)SP = S(S - (b+2)P) = S(S+2P),$$

then we must have $-b+2=2$, i.e., $b=-4$ and then $a=7$, which gives exactly the original inequality.

But even though, for other values of a and b , we are not able to obtain precisely the expression $S+2P$, i.e., a perfect square, we still find interesting and not trivial inequalities.

1. If we take $a=5$, then by (7) we get $b=-3$ and obtain the following new inequality:

Problem 2. Prove that the inequality:

$$\frac{\sqrt{x^2+xy+y^2}}{x^2-xy+y^2} \leq \frac{\sqrt{6}}{\sqrt{x^2+y^2}}, \quad (8)$$

holds for all real numbers x and y , not both equal to 0.

2. If we take $a=9$, then by (7) we get $b=-5$ and obtain the following new inequality:

Problem 3. Prove that the inequality:

$$\frac{\sqrt{x^2+3xy+y^2}}{x^2-xy+y^2} \leq \frac{\sqrt{10}}{\sqrt{x^2+y^2}}, \quad (9)$$

holds for all real numbers x and y , not both equal to 0.

3. If we take $a=11$, then by (7) we get $b=-6$ and obtain the following new inequality:

Problem 4. Prove that the inequality:

$$\frac{\sqrt{x^2+4xy+y^2}}{x^2-xy+y^2} \leq \frac{2\sqrt{3}}{\sqrt{x^2+y^2}}, \quad (10)$$

holds for all real numbers x and y , not both equal to 0.

4. If we take $a=15$, then by (7) we get $b=-8$ and obtain the following new inequality:

Problem 5. Prove that the inequality:

$$\frac{\sqrt{x^2 + 6xy + y^2}}{x^2 - xy + y^2} \leq \frac{4}{\sqrt{x^2 + y^2}}, \quad (11)$$

holds for all real numbers x and y , not both equal to 0.

In the end of this note, notice that there is an essential difference between the first and the second solution of Problem 1: while the later opened a door for further investigations, the former did not.

This is the reason why we can call a solution like the second one, as a creative solution, see [1].

Acknowledgements. During the second part of February 2010, I paid a short visit to University of Rousse, Bulgaria. My hosts there have been my old friends *Emilia Velikova* and *Slavy Bilchev*. At that time Professor *Svetoslav Bilchev* has been very seriously ill and one month later, at the end of March, he passed away. Despite of his illness, he insisted to meet each other and thus we have had our last three unforgettable dinners.

This is a modest homage to *Slavy's* special interest in *Gazeta Matematică*, a journal which he regularly read and used in his scientific and teaching activity.

REFERENCES

- [1] Berinde, V., *Exploring, Investigating and Discovering in Mathematics*, Birkhäuser, Basel, 2004.
- [2] Bilchev, S. I., *Mladejka Balkanska Olimpiada po Matematika 1997-2009*, Mediatex-Pleven, Rousse, 2009.
- [3] Brânzei, D., Șerdean I., Șerdean V., *Junior Balkan Mathematical Olympiad*, Editura Plus, 2003.

GENERALIZĂRI ALE UNOR FORMULE TRIGONOMETRICE

MARIN TOLOȘI¹⁾ și MARIA ALECU²⁾

Abstract. This article establishes formulae for the sine, the cosine and the tangent of a sum of several reals and gives a sufficient condition for their validity

Keywords: domain of the tangent and applications

MSC : 26A09

În prima parte a acestei note, ne-am propus să generalizăm formulele de calcul a cosinusului și sinusului sumei a două numere reale, precum și formulele de transformare a produsului de două cosinusuri, respectiv două sinusuri în sume.

¹⁾Profesor, Colegiul Național „Radu Greceanu“, Slatina.

²⁾Elevă, Colegiul Național „Radu Greceanu“, Slatina.

SUMARUL

Articole și Note matematice

1. Asupra unei probleme deschise din G. M.-B nr. 10/2008, de Árpád Bényi și Cosmin Nițu 337
2. On a homogeneous inequality given at the 2004 J.B.M.O., de Vasile Berinde .. 340
3. Generalizări ale unor formule trigonometrice, de Marin Toloși și Maria Alecu 343
4. Rangul sumei și produsului a două matrice, de Vasile Pop 351
5. Aplicații la inegalitatea lui J. Radon, de D. M. Băținețu-Giurgiu 359
6. Aplicații ale cazului de congruență L.L.U., de Dorel Miheț 362
7. Asupra unor inegalități condiționate, de Dorel Băițan și Dan-Ștefan Marinescu 365

Examene și Concursuri

8. Concursul de matematică „Danubius”, Ediția a IV-a, 24 aprilie 2010, Corabia, Olt, prezentare de Dorin Mărghidanu 370
9. Concursul de matematică „Speranțe”, Ediția a VI-a, 23-25 aprilie 2010, Comănești 375
10. Concursul de Matematică „Dimitrie Pompeiu”, Ediția a X-a, Botoșani, 14 - 16 mai 2010, prezentare de Mircea Trifu 379
11. Concursul de Matematică „N. N. Mihăileanu”, Ediția a XI-a, Constanța, 28-29 mai 2010, prezentare de Gabriela Constantinescu și Cătălin Zîrnă ... 386

Probleme

12. Rezolvarea problemelor din Gazeta Matematică nr. 1/2010 393
13. Rezolvarea problemelor din Gazeta Matematică nr. 2/2010 406
14. Rezolvarea problemelor din Gazeta Matematică nr. 3/2010 418
15. Probleme pentru ciclul primar (P:278 – P:292) 430
16. Probleme pentru gimnaziu (E:14031 – E:14062) 431
17. Probleme pentru liceu (26327 – 26354) 435
18. Probleme pentru concursuri și olimpiade (C.O:5131 – C.O:5146) 439
19. Problems for competitions and olympiads (C.O:5131 – C.O:5146) 441

In memoriam

20. Profesorul Niculae Soare (1949-2009), de Liviu Nicolescu și Gabriel Teodor Pripoae 443

Rubrica rezolvitorilor de probleme

© Toate drepturile privind reproducerea, parțială sau totală, sub orice formă, a materialelor publicate în Gazeta Matematică, sunt rezervate Societății de Științe Matematice din România.



Colecția completă a Gazetei Matematice seria B, în format electronic este realizată de Intuitext (www.intuitext.ro) în parteneriat cu Societatea de Științe Matematice din România.

Preț 15 lei