

## ITERATIVE APPROXIMATION OF FIXED POINTS FOR PSEUDO-CONTRACTIVE OPERATORS

VASILE BERINDE

Department of Mathematics and Computer Science  
Faculty of Sciences North University of Baia Mare

**Abstract.** For a nonexpansive (Lipschitzian) operator  $F$  which is not a strict contraction, the Picard iteration does not converge generally to a fixed point. By adding some pseudo-contractive type assumptions, it is possible to show that several other iterations (Krasnoselski-Schaefer, Mann, Ishikawa) converge to a fixed point of  $F$ .

The main aim of the paper is to survey some old and recent results especially related to the convergence of Krasnoselski-Schaefer iteration in the class of generalized pseudo-contractive and Lipschitzian operators.

### 1. INTRODUCTION

Many of the most important nonlinear problems of applied mathematics reduce to finding solutions of nonlinear functional equations (integral equations, differential equations etc.) which can be formulated in terms of finding the fixed points of a given nonlinear operator of an infinite dimensional function space  $X$  into itself:

$$(1) \quad x = Tx$$

There is a classical general existence theory of fixed points for mappings satisfying compactness conditions associated with the names of Brower, Schauder, Leray etc. as well as an abundant literature of metrical fixed point theorems, which establish the existence (and uniqueness) of fixed points for maps satisfying a variety of contractive conditions ([16]). The first basic result is the classical Picard-Banach-Caccioppoli principle

**THEOREM 0.** *Let  $(X, d)$  be a complete metric space and  $T : X \rightarrow X$  a strict contraction, that is, there exists  $\alpha, 0 < \alpha < 1$  such that*

$$(2) \quad d(Tx, Ty) \leq \alpha \cdot d(x, y) \text{ for all } x, y \in X$$

*Then the Picard iteration (the sequence of successive approximation)  $(x_n)$ , given by*

$$(3) \quad x_n = T(x_{n-1}) = T^n(x_0), \quad n = 1, 2, \dots$$

*converges to the unique fixed point  $x^*$  of  $T$ ,*

$$x_n \rightarrow x^* \text{ (as } n \rightarrow \infty \text{)}.$$

The convergence order of the Picard iteration in Theorem 0 is given by both *a priori* and *a posteriori* estimates:

$$(4) \quad d(x_n, x^*) \leq \frac{\alpha^n}{1-\alpha} \cdot d(x_0, x_1), \quad n \geq 1;$$

$$(4') \quad d(x_n, x^*) \leq \frac{\alpha}{1-\alpha} \cdot d(x_n, x_{n-1}), \quad n \geq 1;$$

This fundamental result in the fixed point theory has been extended to some larger classes of contractive and generalized contractive operators, see [16], for example, by replacing the strict contractive condition (2) by a weaker condition of the following type

$$(2') \quad d(Tx, Ty) \leq \varphi(d(x, y)), \quad x, y \in X$$

where  $\varphi: \mathbb{R}_+ \rightarrow \mathbb{R}_+$  is a certain comparison function or by a more general one

$$d(Tx, Ty) \leq \varphi(d(x, y), d(x, Tx), d(x, Ty), d(y, Tx), d(y, Ty)), \quad x, y \in X,$$

where  $\varphi: \mathbb{R}_+^5 \rightarrow \mathbb{R}_+$  stands for a certain 5-dimensional comparison function (see [2], [21]). Important results as Kannan or Ćirić theorems belong to the second class of generalized contractions [16].

If the generalized contractive condition in a such fixed point theorem is strong enough, that is, the comparison function satisfies some essential conditions, then the (possible unique) fixed point of  $T$  can be obtained by means of the sequence of successive approximations (Picard iteration).

But, if the contractive condition is slightly weaker, as for example the case when  $T$  is only nonexpansive

$$(2'') \quad d(Tx, Ty) \leq d(x, y), \quad x, y \in X,$$

then the Picard iteration (3) need no longer converge to a fixed point of  $T$  (if any).

In fact, in general, a nonexpansive operator  $T$  need not have a fixed point and even if  $T$  possesses a fixed point, the Picard iteration need not converge to this point, see Example 1 in Section 2.

Even if the fixed point can be obtained by the Picard iteration, it is of interest to determine other iteration procedures that could converge faster, in a certain sense, to the fixed point.

## 2. OTHER FIXED POINT ITERATION PROCEDURES

The next example illustrates the case of a nonexpansive operator possessing an unique fixed point, for which the Picard iteration does not converge to that fixed point, except for the case when the initial approximation coincides to the fixed point itself.

**EXAMPLE 1.** ([9]) Let  $X = [0, 1]$  and  $T: [0, 1] \rightarrow [0, 1]$  be the linear function

$$T(x) = 1 - x, \quad 0 \leq x \leq 1.$$

- Then :    a)  $T$  is nonexpansive;  
           b)  $T$  has an unique fixed point

$$F_T = \{x \in [0, 1] / T(x) = x\} = \left\{\frac{1}{2}\right\}.$$

c) The Picard iteration  $x_n = T(x_{n-1}), n = 1, 2, \dots$ , yields the oscillatory sequence  $a, 1 - a, a, 1 - a, a, \dots$  for any  $x_0 = a (a \neq \frac{1}{2})$ .

Taking into account the fact that the class of nonexpansive operators is very important in applications, we have to impose certain additional conditions on the ambient space or on the operator itself, in order to ensure the existence of a fixed point or to guarantee the convergence of the Picard iteration to a fixed point of the operator.

In what concern the existence problem, the following result was obtained independently by Browder, Gohde and Kirk (see [3], for example):

**THEOREM 1.** *Let  $C$  be a closed, bounded, and convex subset of a uniformly convex Banach space,  $T : C \rightarrow C$  a nonexpansive map. Then  $T$  has a fixed point.*

The proofs of Theorem 1 are, unfortunately, all not constructive. As shown by Example 1, the Picard iteration does not converge (to the fixed point) of such a nonexpansive operator.

A similar situation is encountered for the class of lipschitzian (and pseudo-contractive, in some sense) operators, when, even if the fixed point is unique, the Picard iteration does not converge (see [1]).

To remove these difficulties we need to consider some other sequential procedures to be used for approximating fixed points.

We present here a few chronological reference points. In 1953 W. R. Mann [17] introduced an iteration procedure which can be represented in the following form

$$(5) \quad x_{n+1} = (1 - a_n)x_n + a_n T x_n, \quad n = 0, 1, 2, \dots$$

where  $(a_n)$  is sequence satisfying: (i)  $a_n \in [0, 1]$  and (ii)  $\sum a_n = \infty$ .

Mann showed that, if  $T$  is an continuous selfmap of a closed interval  $[a, b]$  with at most one fixed point, then the iteration scheme (5), with  $a_n = 1/(n + 1)$ , converges to the fixed point of  $T$ .

In 1955, Krasnoselski [18] showed that, if  $X$  is a uniformly convex Banach space, and  $T : X \rightarrow X$  is nonexpansive, then the iteration  $(x_n)$ ,

$$(6) \quad x_{n+1} = \frac{1}{2}(x_n + T x_n), \quad n = 0, 1, 2, \dots$$

converges to a fixed point of  $T$ .

Later, in 1957, Schaefer considered the extension of (6), by replacing  $1/2$  by a constant  $\lambda \in [0, 1]$ , that is, he introduced the iteration procedure  $(x_n)$ ,

$$(6') \quad x_{n+1} = (1 - \lambda)x_n + \lambda \cdot T x_n, \quad n = 0, 1, 2, \dots$$

and proved similar results [19].

It is easy to see that  $(x_n)$  given by (6') is in fact the Picard iteration for the associated operator  $U = (1 - \lambda) \cdot I + \lambda \cdot T$ , where  $I$  is the identity.

If we take  $\lambda = \frac{1}{2}$ , from (6') we obtain (6), and, by putting  $\lambda = 1$ , from (6') we obtain the Picard iteration. Iteration (6'), which will be called in the sequel *Krasnoselski-Schaefer iteration*, is a particular case of the *Mann iteration* (5), obtained from the last one for  $a_n = \lambda$  (*const*).

It was proven for continuous mappings that, if the Mann iterative process converges, then it must converge to a fixed point of  $T$ . But if  $T$  is not continuous, there is no guarantee that, even if the Mann process converges, it will converge to a fixed point of  $T$ , as shown by the next example.

**EXAMPLE 2.** Let  $X = [0, 1]$  and  $T : X \rightarrow X$  the map defined by  $T0 = T1 = 0$  and  $Tx = 1$ , for  $x \in (0, 1)$ .

Then  $T$  is a selfmap of  $[0, 1]$ , having the unique fixed point  $x = 0$ . However, the Mann iteration, with  $a_n = 1/(n+1)$  and  $x_0 \in (0, 1)$  converges to 1, which is not a fixed point of  $T$ .

By adding some pseudocontractive hypotheses to a nonexpansive map, Ishikawa [7] proved the following result.

**THEOREM 3.** Let  $E$  be a convex, compact subset of a Hilbert space  $H$ ,  $T : E \rightarrow E$  a Lipschitzian and pseudocontractive map. Then the sequence  $(x_n)$ , defined by

$$(7) \quad x_{n+1} = (1 - \alpha_n) \cdot x_n + \alpha_n T [(1 - \beta_n)x_n + \beta_n T x_n], \quad n = 0, 1, 2, \dots$$

where  $(\alpha_n), (\beta_n)$  are sequences of positive numbers satisfying the conditions  $0 \leq \alpha_n \leq \beta_n \leq 1$ ,  $\lim \beta_n = 0$  and  $\sum \alpha_n \beta_n = \infty$ , converges strongly to a fixed point of  $T$ .

The next diagram represents the most important iteration procedures considered here

$$x_{n+1} = Tx_n, \quad n \geq 0 \\ \uparrow \lambda = 1$$

1890 Picard

$$x_{n+1} = \frac{1}{2}(x_n + Tx_n), \quad n \geq 0 \\ \uparrow \lambda = \frac{1}{2}$$

1955, Krasnoselski

$$x_{n+1} = (1 - \lambda)x_n + \lambda Tx_n, \quad n \geq 0, 0 < \lambda \leq 1 \\ \uparrow a_n = \lambda(\text{const})$$

1957 (Krasnoselski-)Schaeffer

$$x_{n+1} = (1 - a_n) \cdot x_n + a_n \cdot Tx_n, \quad n \geq 0, a_n \in [0, 1] \\ \uparrow b_n = 0$$

1953 Mann

$$x_{n+1} = (1 - a_n) \cdot x_n + a_n T [(1 - b_n)x_n + b_n T x_n], \quad n \geq 0, 0 \leq a_n, b_n \leq 1$$

1974 Ishikawa

## 3. PSEUDO-CONTRACTIVE OPERATORS

The interest in **pseudocontractive mappings** is due mainly to

- a) their usefulness as an additional assumption to Lipschitz type conditions in proving convergence of fixed point iterative procedures;  
 b) their connection with the important class of nonlinear **accretive operators**.

Let  $H$  be a Hilbert space and  $T : H \rightarrow H$  a selfmap.

**Definition 1.**  $T$  is said to be **pseudoccontractive** on  $C \subset H$  if

$$(10) \quad \|Tx - Ty\| \leq \|x - y\|^2 + \|Tx - Ty - (x - y)\|^2, \quad x, y \in C$$

or equivalently

$$(10') \quad \langle Tx - Ty, x - y \rangle \leq \|x - y\|^2, \quad x, y \in C.$$

**Definition 2.**  $T$  is called **generalized pseudocontractive**, if  $\exists r > 0$  such that

$$(11) \quad \|Tx - Ty\|^2 \leq r \cdot \|x - y\|^2 + \|Tx - Ty - r(x - y)\|^2, \quad x, y \in C,$$

or equivalently

$$(11') \quad \langle Tx - Ty, x - y \rangle \leq r \cdot \|x - y\|^2, \quad x, y \in C$$

**Remark.** For  $r = 1$ , from Definition 2 we obtain Definition 1.

**Definition 3.** The operator  $T$  is said to be **strictly pseudocontractive on C** if there exists  $k < 1$  such that

$$(12) \quad \|Tx - Ty\|^2 \leq \|x - y\|^2 + k \|Tx - Ty - (x - y)\|^2, \quad x, y \in C,$$

**Definition 4.** The operator  $T$  is called **strongly pseudocontractive on C** if there exist  $t > 1$  and  $r > 0$  such that

$$\|x - y\|^2 \leq \|(1 + r)(x - y) - rt(Tx - Ty)\|, \quad x, y \in C.$$

**Remarks.** 1) For  $t = 1$ , from Definition 4 we obtain Definition 1;

2)  $T$  is pseudo-contractive if and only if  $I - T$  is accretive ([21]).

The class of pseudo-contractive operators usually associated with lipschitzian properties has been studied extensively by various authors, see [22] and references therein.

The following result has been proved by Verma [12].

**THEOREM 4.** Let  $K$  be a non-empty closed convex subset of  $H$  and  $T : K \rightarrow K$  a lipschitzian and generalized pseudocontractive operator (with constant  $s$  and  $r$ , respectively,  $r < 1$ ).

Then, for any  $\lambda, 0 < \lambda < \frac{2(1-r)}{1-2r+s^2}$  the iteration  $(x_n)$ , given by

$$x_{n+1} = (1 - \lambda)x_n + \lambda Tx_n, \quad n \geq 0$$

converges strongly to the unique fixed point  $x^*$  of  $T$ :

$$x_n \rightarrow x^* (n \rightarrow \infty).$$

In our paper [1] we completed the result of Verma by inserting both *a priori* and *a posteriori* error estimates (Theorem 3.1) and by finding the fastest iteration amongst all Krasnoselski-Schafer iterations.

Moreover, when the Picard iteration and Krasnoselski-Schafer iteration converge simultaneously, it is possible to compare the fastest one, in a certain sense of rate of convergence. Generally, it is possible to find the fastest Krasnoselski-Schafer iteration in the family, as shown by

**THEOREM 5** ([1]). *Let  $K$  be a non-empty closed convex subset of a Hilbert space and  $T : K \rightarrow K$  a generalized pseudocontractive (with constant  $r$ ,  $0 < r < 1$ ) and Lipschitzian (with constant  $s > 0$ ) operator and  $\lambda \in (0, 1)$  such that  $0 < \lambda < \frac{2(1-r)}{1-2r+s^2}$ .*

Then

- (i)  $T$  has an unique fixed point  $x^* \in K$ ;  
 (ii) The Krasnoselski-Schafer iteration  $(x_n)$  converges strongly to  $x^*$ , for each  $x_0 \in K$ ;

$$(iii) \quad \|x_n - x^*\| \leq \frac{\theta^n}{1-\theta} \cdot \|x_1 - x_0\|, \quad n \geq 1$$

$$\|x_n - x^*\| \leq \frac{\theta}{1-\theta} \cdot \|x_n - x_{n-1}\|, \quad n \geq 1$$

where  $\theta = \sqrt{(1-\lambda)^2 + 2\lambda(1-\lambda)r + \lambda^2 s^2}$ .

- (iv) The fastest Krasnoselski-Schafer iteration is obtained for

$$\theta_0 = \frac{1-r}{1-2r+s^2}.$$

**EXAMPLE 3.**([1]) Let  $K = [\frac{1}{2}, 2]$  and  $T : K \rightarrow K$  the function defined by  $T(x) = \frac{1}{x}$ ,  $x \in [\frac{1}{2}, 2]$ . Then

- 1)  $T$  is Lipschitzian with constant  $s = 4$ ;
- 2)  $T$  is generalized pseudocontractive with constant  $r > 0$  arbitrary;
- 3) The Picard iteration, with  $x_0 = a \neq 1$  yields the oscillatory sequence

$$a, \frac{1}{a}, a, \frac{1}{a}, \dots$$

- 4) The Krasnoselski-Schafer iteration converges to  $x^* = 1$ , for any  $\lambda \in (0, \frac{2(1-r)}{1-2r})$ ,  $r < 1$ ;

- 5) For  $r = 0,5$ , the fastest Krasnoselski-Schafer iteration is obtained taking  $\lambda_0 = \frac{1}{32}$ , i.e.

$$x_{n+1} = \frac{1}{32} \left( 31x_n + \frac{1}{x_n} \right), \quad n \geq 0$$

which converges to  $x^* = 1$ , (slowly, because the contraction coefficient  $\theta_0 = \frac{\sqrt{63}}{8} \approx 0,992$  is very close to 1).

#### FINAL REMARKS

The class of pseudo-contractive operators has been intensively studied in the last decade (see, for example, [4], [5]). We considered a more general class of pseudo-contractive operators in [20].

## REFERENCES

- [1] BERINDE, V., The fastest iteration for approximating fixed points of lipschitzian pseudo-contractions, *Proceed. 2nd Palestinian Int. Math. Conference* (in print).
- [2] BERINDE, V., *Generalized contractions and applications* (in Romanian), Editura CUB PRESS 22, Baia Mare, 1997.
- [3] BROWDER, F.E., PETRYSHIN, W.V., Construction of fixed points of nonlinear mappings in Hilbert spaces, *J. Math. Anal. Appl.* 20(1967), 197-228.
- [4] CHIDUME, C.E., Global iteration schemes for strongly pseudo-contractive maps, *Proc. Amer. Mat. Soc.* 126(1998), 9, 2641-2649.
- [5] CHIDUME, C.E., Approximation of fixed points of strongly pseudocontractive mappings, *Proc. Amer. Mat. Soc.* 120(1994), 2, 515-551.
- [6] GHOSH, M.K., DEBNATH, L., Convergence of Ishikawa iterates of quasi-nonexpansive mappings, *J. Math. Anal. Appl.* 207(1997), 96-103.
- [7] ISHIKAWA, S., Fixed points by a new iteration method, *Proc. Amer. Mat. Soc.* 44(1974), 1, 147-150.
- [8] PETRYSHYN, W.V., WILLIAMSON, T.E., Strong and weak Convergence of the Sequence of Successive Approximations for Quasi-Nonexpansive Mappings, *J. Math. Anal. Appl.* 43(1973), 459-497.
- [9] RHOADES, B.E., Some fixed point iteration procedure, *Internat. J. Math.&Math.Sci.*, 14(1991), 1, 1-16.
- [10] RHOADES, B.E., A general principle for Mann iterations, *Indian. J. Pure Appl. Math.* 26(8), 1995, 751-762.
- [11] RHOADES, B.E., Convergence of an Ishikawa-Type Iteration Scheme for a Generalized Contraction, *J. Math.Anal. Appl.*, 185(1994), No. 2, 350-355.
- [12] VERMA, R.U., A fixed point theorem involving lipschitzian generalized pseudo-contractions, *Proceed. Royal Irish.*, 97A(1997), No.1, 83-86.
- [13] VERMA, R.U., Strongly nonlinear variational inequalities and generalized pseudocontractions, *Rend. Circ. Mat. Palermo*, XLVIII(1999), 201-208.
- [14] BERINDE, V., On some stability results for fixed point iterative procedures, (submitted).
- [15] PICARD, É., Mémoire sur la théorie des equations aux dérivée partielles et la metode des approximations successives, *J. Math. Pures. Appl.*, 6(1890), 145-210.
- [16] RUS, A.I., *Generalized contractions and applications*, Cluj University Press, Cluj-Napoca, 2001.
- [17] MANN, W.R., Mean value methods in iteration, *Proc. Amer. Mat. Soc.* 4(1953), 506-510.
- [18] KRASNOSELSKI, M.A., Two remarks on the method of successive approximations, *Uspechi Mat. Nauk.* 10(1955), No. 1(63), 123-127.
- [19] SCHAEFER, H., Uber die Methode Sukzessiver Approximation, *Jber. Deutsch Math.-Verein*, 59(1957), 131-140.
- [20] BERINDE, V., On the convergence of Krasnoselski-Schaefer iteration for pseudo- $\varphi$ -contractions (in preparation).
- [21] BROWDER, F.E., Nonlinear mappings of nonexpansive and accretive type in Banach spaces, *Bull. Amer. Math. Soc.* 73(1967), 875-882.
- [22] CHIDUME, C.E., Global iteration schemes for strongly pseudocontractive maps, *Proc. Math. Soc.* 126(1998), 2641-2649.